



Environment

Freshwater: Yes/ No

Habitat: Marine

Salinity:

Brackish: Yes/No


Migrations:

Temperature:

Salt Water: Yes✓/ No

Depth range :

Picture (scanned images or photographs of adult/ larval stages)



*Lanceola loveni antarctica* Vinogradov,  
female (after Vinogradov,1962)

|  |  |  |  |
|--|--|--|--|
| <p>DATA ENTRY FORM: Form -2 (Fish/ Shell fish/ Others ) Ref. No.:</p> <p>(Please answer only relevant fields; add additional fields if you require)</p> <p>Form- 1 Ref. No.:</p>   |  |  |  |
| <p>IMPORTANCE</p> <p>Landing statistics (t/y): from                      to                      Place:                      Ref. No.:</p> <p>Main source of landing: Yes/ No                      Coast: east/ west</p> <p>Importance to fisheries:</p> <p>Main catching method:</p> <p>Used for aquaculture: yes/ never/ rarely</p> <p>Used as bait: yes/no/ occasionally</p> <p>Aquarium fish: yes/ no/ rarely</p> <p>Game fish: yes/ no</p> <p>Dangerous fish: poisonous/ harmful/ harmless</p> <p>Bioactivity: locally known/ reported/ not known                      Details:</p> <p>Period of availability: Throughout the year – yes/ no                      If no, months:</p>  |  |  |  |
| <p>SALIENT FEATURES:</p> <p>Morphological:</p> <p>Diagnostic characteristics:</p> <p>Individuals of all ages have dorsal denticles on the pleon somites and denticles on somite I of the urosoma which are stouter than in the typical form, the 6<sup>th</sup> segment I of the urosoma which are rami of the uropods are relatively longer and narrower. The two latter differences are well marked in young specimens (7-12 mm long) and are practically absent in larger individuals (19-27) mm long). True, even in the typical individuals the young specimens have a narrower 6<sup>th</sup> segment in pereopods I and relatively longer rami in the uropods than do the adults. However, in them the age-dependent changes are not manifest as sharply as in <i>L. loveni antarctica</i>.</p> |  |  |  |
| <p>Sex attributes:</p> <p>Dimorphic</p> <p>Male: 1<sup>st</sup> antenna well developed , female: 1<sup>st</sup> antenna reduced.</p> <p>Descriptive characters:</p>  |  |  |  |

Meristic characteristics:

Feeding habit:

Main food:

Feeding type:

Additional remarks:

Size and age:

Maximum length (cm) (male/ female/ unsexed)

Ref. No.:

Length of sexually mature individuals up to 27mm.

Average length (cm) (male/female/unsexed)

Ref. No.:

Maximum weight: (g) (male/female/unsexed)

Ref. No.:

Average weight: (g) (male/female/unsexed)

Ref. No.:

Longevity (y) (wild): (captivity)

Ref. No.:

Length/ weight relation ships:

|  |  |
|--|--|
| Eggs and larvae:<br>Characteristics:<br>Abundance:<br>Biochemical aspects:<br>Proximate analysis: moisture/ fat/ protein/ carbohydrate/ash<br>Electrophoresis:   | Ref. No.<br><br><br><br><br>Ref. No.<br>Ref. No. |
| <b>SPAWNING INFORMATION:</b><br>Locality:<br>Season:<br>Fecundity:<br>Comment:   | Main Ref:  |
| <b>MAJOR PUBLICATIONS (INDIAN):</b><br>(Include review articles, monographs, books etc.)<br><br><b>LIST OF INDIAN EXPERTS (Name, address, phone, fax, e-mail etc.)</b><br><br><div style="margin-left: 40px;"> <p>Dr.K.K.C.Nair<br/>           Scientist-In-Charge<br/>           R.C. of NIO,<br/>           Post Box-1616<br/>           Kochi – 682 014</p> <p>Dr. N. Krishna pillai<br/>           “Radhika”<br/>           65- Champaka Nagar<br/>           Bakery Junction<br/>           Trivandrum-695 001</p> </div> |  |
| <b>ACKNOWLEDGMENT:</b><br>(List of persons who contributed, modified or checked information)   |  |